

VOLUME 29

OCTOBER 1951

NUMBER 5

*Canadian
Journal of Zoology*

***Editor:* E. HORNE CRAIGIE**

***Published by* THE NATIONAL RESEARCH COUNCIL
OTTAWA CANADA**

CANADIAN JOURNAL OF ZOOLOGY

This was formerly *Section D, Canadian Journal of Research*. The change to the new name took place January 1, 1951. The CANADIAN JOURNAL OF ZOOLOGY is published six times annually.

The CANADIAN JOURNAL OF ZOOLOGY is published by the National Research Council of Canada under the authority of the Chairman of the Committee of the Privy Council on Scientific and Industrial Research. Matters of general policy are the responsibility of a joint Editorial Board consisting of members of the National Research Council of Canada and the Royal Society of Canada.

The National Research Council of Canada publishes also the *Canadian Journal of Botany*, *Canadian Journal of Chemistry*, *Canadian Journal of Medical Sciences*, *Canadian Journal of Physics*, *Canadian Journal of Technology*.

EDITORIAL BOARD

<i>Representing</i>	<i>Representing</i>	
NATIONAL RESEARCH COUNCIL	ROYAL SOCIETY OF CANADA	
DR. J. H. L. JOHNSTONE, (<i>Chairman</i>), Professor of Physics, Dalhousie University, Halifax, N.S.	DR. G. M. VOLKOFF, Professor of Physics, University of British Columbia, Vancouver, B.C.	} Section III
DR. OTTO MAASS, Macdonald Professor of Physical Chemistry, McGill University, Montreal, P.Q.	DR. J. W. T. SPINKS, Dean, College of Graduate Studies, University of Saskatchewan, Saskatoon, Sask.	
DR. CHARLES W. ARGUE, Dean of Science, University of New Brunswick, Fredericton, N.B.	DR. H. S. JACKSON, Head, Department of Botany, University of Toronto, Toronto, Ont.	
DR. A. G. MCCALLA, Department of Plant Science, University of Alberta, Edmonton, Alta.	DR. E. HORNE CRAIGIE, Department of Zoology, University of Toronto, Toronto, Ont.	} Section V
<i>Ex officio</i>	<i>Representing</i>	
DR. LÉO MARION, Editor-in-Chief, Division of Chemistry, National Research Laboratories, Ottawa.	THE CHEMICAL INSTITUTE OF CANADA	
DR. H. H. SAUNDERSON, Director, Division of Information Services, National Research Council, Ottawa.	DR. H. G. THODE, Department of Chemistry, McMaster University, Hamilton, Ont.	

Subscription rate: \$2.00 a year. All enquiries concerning subscriptions should be addressed to the CANADIAN JOURNAL OF ZOOLOGY, National Research Council, Ottawa, Canada. Special rates can be obtained for subscriptions to more than one of the Journals published by the National Research Council.

Canadian Journal of Zoology

Issued by THE NATIONAL RESEARCH COUNCIL OF CANADA

VOL. 29

OCTOBER, 1951

NUMBER 5

THE EFFECTS ON THE GRASSHOPPER, *MELANOPLUS MEXICANUS MEXICANUS* (SAUSS.) (ORTHOPTERA: ACRIDIDAE), OF VARYING THE NITROGEN CONTENT IN ITS FOOD PLANT¹

BY D. S. SMITH² AND F. E. NORTHCOTT³

Abstract

Melanoplus mexicanus mexicanus (Sauss.), from time of hatching, was fed on wheat (Renown) grown in nutrient solutions of varying nitrogen content. Some wheat was produced with an average nitrogen content of 6.16% (dry weight), some with 4.29%, and some with 3.33%. Survival and development were greatest on the high nitrogen wheat, least on the low. On the low nitrogen wheat no individuals developed beyond the last nymphal instar. Neither weight of adults nor the sex ratio was affected by the various foods. Females fed on wheat with a high nitrogen content laid more eggs but the viability of the eggs was not affected.

Introduction

Insects are known to differ in such biological factors as survival, rate of development, and number of progeny, depending upon the food they eat. In phytophagous insects these differences may be caused by feeding on plants of different genera, of different species, or even of different varieties of a species. Among innumerable papers on this might be mentioned those on Orthoptera by Hodge (11), Chauvin (3), Smith, Handford, and Chefurka (19), and Tauber, Drake, and Decker (20); those on Lepidoptera by Basu (2) and Seamans and McMillan (18); and one on Coleoptera by Hill (10).

Some attempts have been made to determine the contributing factors. Webster, Seiglinger, and Davies (21) made extensive chemical analyses of a number of varieties of sorghum but found no substances that could be correlated with the differences in longevity and fecundity found by Dahms (5) in chinch bugs feeding on these varieties. Creighton (4) stated that *Alabama argillacea* (Hbn.) underwent a prolonged developmental period and suffered a very high mortality rate on zinc- and copper-deficient plants (presumably cotton). Pepper and Hastings (17) suggest that sterility in the sugar-beet webworm is connected with an insufficient amount of linoleic acid in the host plant.

¹ Manuscript received May 15, 1951.

Contribution No. 2832, Division of Entomology, Science Service, Department of Agriculture, Ottawa, Canada.

² Agricultural Research Officer, Field Crop Insect Laboratory, Lethbridge, Alta.

³ Technical Officer, Field Crop Insect Laboratory, Lethbridge, Alta.

[The August issue of this Journal (Can. J. Zool. 29: 265-295. 1951) was issued Aug. 13, 1951.]

Much of this work has been concerned with the amount of nitrogen in the food plant. Mumford and Hey (16) claim that, in general, a plant with a high nitrogen content stimulates reproduction in insects feeding on it. Evans (8) has correlated the rate of reproduction of an aphid, *Brevicoryne brassicae* (L.), with the nitrogen content of the host. Haseman (9) reported that the chinch bug would mature faster and produce more offspring on nitrogen-deficient corn. On the other hand he found that *Toxoptera graminum* (Rond.) produced very few offspring on nitrogen-deficient plants.

This question of the factors in the host plant that affect the insect attacking it is important in developing varieties of plants resistant to insects or unsuitable for them. In some cases resistance may depend on such a nutritional factor (15). Dahms (5) has shown that the chinch bug survives better and has a greater fecundity on susceptible varieties of sorghum, and Davidson (6) has had similar results with aphids on beans. It has been suggested that the nitrogen content of the cell sap of plants is an important factor in the susceptibility of the plant to insect attack (1, 7, 12).

The present study investigates the effect of the amount of nitrogen in a food plant on the lesser migratory grasshopper, *Melanoplus mexicanus mexicanus* (Sauss.). Wheat was grown in nutrient solutions to produce plants containing various amounts of nitrogen. These plants were used as the sole food for the grasshopper during the whole of its nymphal and adult life.

Materials and Methods

Renown wheat was grown outdoors in asphalt-coated basins filled with vermiculite (Terralite). Seeds were planted in this material and watered with distilled water until they sprouted. Then nutrient solutions prepared with distilled water were supplied to the basins every six hours by an automatic subirrigation method adapted from one described by Lott (13). Three nutrient solutions were used: high nitrogen, low nitrogen, zero nitrogen (Table I).

TABLE I
COMPOSITIONS OF NUTRIENT SOLUTIONS FOR WHEAT

Stock solution		Zero nitrogen, ml./liter	Low nitrogen, ml./liter	High nitrogen, ml./liter
KH ₂ PO ₄	1 M	—	6.45	—
Ca(NO ₃) ₂	1 M	—	0.25	31.50
MgSO ₄	1 M	2.00	2.00	2.00
(NH ₄) ₂ SO ₄	1 M	—	0.10	—
CaCl ₂	1 M	—	2.75	—
NH ₄ H ₂ PO ₄	1 M	—	—	6.45
NH ₄ NO ₃	1 M	—	—	33.50
KNO ₃	1 M	—	—	6.45
K ₂ SO ₄	0.25 M	10.00	—	—
Ca(H ₂ PO ₄) ₂	0.05 M	40.00	—	—
CaSO ₄	0.01 M	200.00	—	—

To each liter of these solutions was added one milliliter of a stock solution of micronutrient elements prepared as follows:—

<i>Ingredient</i>	<i>Grams per liter</i>
H ₃ BO ₃	2.86
MnCl ₂ · 4H ₂ O	1.81
ZnSO ₄ · 7H ₂ O	0.22
CuSO ₄ · 5H ₂ O	0.08

Ferrous sulphate was applied to the surface of the vermiculite in the basins at the rate of 10 to 15 p.p.m. of nutrient solution at least once a week. The solutions were completely replaced every two weeks. Weekly measurements of the pH of the solutions showed a range of 4.8 to 6.6, so that no adjustment was considered necessary.

Total nitrogen determinations were made by the method of Ma and Zuazaga (14). Several samples were taken from the wheat grown in each nutrient solution at intervals from 10 to 30 days after germination, the period during which the wheat was used as food. The samples were dried at room temperature, ground, and then further dried to constant weight in an oven at 98° C.

Field-collected eggs of *M. m. mexicanus* were hatched in the laboratory and the nymphs were placed in two-quart sealers. The glass top of the sealer was replaced by wire mesh screening, and an inch of soil was packed in the bottom. The sealers were kept outdoors and shielded with boards from rain and the heat of the midday sun.

Ten sealers, each containing 10 nymphs, comprised one series, and one series was devoted to each type of food. On June 11 three series were started and they are designated below as High N, Low N, and Zero N_a. The mortality rate in the Zero N_a series increased so rapidly that two additional series were started, Zero N_b on July 22, and Zero N_c on Aug. 1. Cuttings from the wheat from the different treatments were supplied to the grasshoppers two to four times daily. The sealers were examined daily and molts and deaths recorded. Each series continued until there was no further oviposition and most of the insects had died.

The soil in the sealers was sifted every few days and the egg pods were removed to cold storage. After about four months at 35°–40° F. they were placed at room temperature. The numbers hatching were recorded as were also the number of eggs that did not hatch.

Nitrogen Content of Wheat

The ranges of total nitrogen content of wheat from the high nitrogen and low nitrogen treatments are quite distinct (Table II). There is, however, an overlapping between that from the low nitrogen treatment and that from the zero.

TABLE II

TOTAL NITROGEN CONTENTS OF WHEAT GROWN IN THREE NUTRIENT SOLUTIONS AS PERCENTAGES OF DRY WEIGHTS

Nutrient solution	Number of determinations	Nitrogen content	
		Range	Average
High nitrogen	42	7.41-5.07	6.16
Low nitrogen	53	4.62-3.79	4.29
Zero nitrogen	45	4.01-2.29	3.33

Survival

The trend of survival of the grasshoppers in the five series during the whole period of nymphal development is shown in Fig. 1.

The grasshoppers on the High N series consistently showed the highest survival and those on the Zero N series the lowest. In Series Zero N_b all individuals were dead in 15 days and in Zero N_a 96% were dead after 25 days. Mortality did not increase so rapidly on series Zero N_c, nor was the final mortality as great as on the other two. This series differed from all others in that the nymphs were hatched from eggs that were collected from a different location and it was started later in the season.

Development

The average durations of the nymphal stadia of the grasshopper are shown in Table III. Figures are not given for the Zero N series, as so few individuals

TABLE III

RATES OF DEVELOPMENT OF *M. m. mexicanus* ON WHEATS GROWN IN SOLUTIONS OF HIGH AND OF LOW NITROGEN CONTENT

Nutrient series	Average nitrogen content of wheat	Average number of days per instar					Average duration of nymphal period in days
		1st	2nd	3rd	4th	5th	
High N	6.16	12	7	6	8	16	49
Low N	4.29	13	7	8	15	18	61

in these lived past the second instar and none completed development. Development was more rapid in the High N series, particularly in the later instars.




FIG. 1. Trend of survival of *M. m. mexicanus* during period of nymphal development on wheats grown in nutrient solutions of different nitrogen contents.

The number of individuals developing to the adult stage on the high nitrogen wheat was significantly greater than the number that developed on the others (Table IV).

TABLE IV
SURVIVAL OF *M. m. mexicanus* AND NUMBERS OF ADULTS DEVELOPING

Nutrient series	Average nitrogen content of wheat	Number surviving after 66 days	Percentage survival	Number of adults	Percentage adults ^a
High N	6.16	44	48	44	48
Low N	4.29	30	30	29	29
Zero N _a	3.33	4	4	0	0
Zero N _b	3.33	0	0	0	0
Zero N _c	3.33	13	16	0	0

* Least significant difference (5% level), 19.

In the Zero N series no adults appeared and most individuals failed to develop beyond the second instar. Development appeared to be more uniform on the High N series, as 90% of those reaching the adult stage did so within 11 days of each other. In the Low N series 25 days elapsed after the appearance of the first adult before 90% were in this stage.

Weight of Adults

Newly molted adults were weighed individually as they appeared. The average weights (Table V) indicate no differences as a result of the different foods. It is unusual, though, for the weight of the male to exceed that of the female, as it does in the Low N series.

TABLE V
AVERAGE WEIGHTS OF NEWLY EMERGED ADULTS

Nutrient series	Average nitrogen content of wheat	Male	Female
High N	6.16	0.1532 gm.	0.1649 gm.
Low N	4.29	0.1613 gm.	0.1561 gm.

Sex Ratio

The numbers of males and females appearing on the High N and the Low N series (Table VI) did not differ significantly from a 50 : 50 ratio and revealed no differential effect of the food on sex.

TABLE VI
NUMBERS OF MALES AND FEMALES IN SURVIVING ADULTS

Nutrient series	Average nitrogen content of wheat	Male	Female	Total
High N	6.16	25	19	44
Low N	4.29	11	18	29

Fecundity and Egg Viability

The numbers of eggs laid by females reared on wheat from the high nitrogen and the low nitrogen treatments as well as the number that hatched are shown in Table VII. Females fed on wheat from the high nitrogen treatment laid considerably more eggs than those fed on low nitrogen wheat. The relative viability of the eggs was measured by the percentage hatch; and as the two groups of eggs were kept together under identical conditions, the figures are considered to be a reliable estimate. The different foods seem to have no effect on viability, the percentage hatch being approximately the same for the two series.

TABLE VII
EGG PRODUCTION AND PERCENTAGE HATCH

Nutrient series	Average nitrogen content of wheat	Total number of eggs	Number of females	Number of eggs per female	Number of eggs hatched	Percentage hatch
High N	6.16	1094	19	57.6	490	44.8
Low N	4.29	624	18	34.7	292	46.8

Conclusion

This study clearly indicates a positive correlation between an increase in the nitrogen content of the wheat and a greater vitality of *M. m. mexicanus* as measured in terms of survival, development, and egg production. The nitrogen content of the food plant had no effect on egg hatchability.

Acknowledgments

The authors wish to express their appreciation to Dr. R. H. Handford, Dominion Entomological Laboratory, Kamloops, B.C., for his encouragement and many helpful suggestions. Mr. G. Thomas, formerly of the Dominion Entomological Laboratory, Brandon, Man., where this project was carried out, and Mr. W. Romanow, also of that laboratory, helped considerably with the rearing.

References

1. AUCLAIR, J. L. and MALTAIS, J. B. Studies on the resistance of plants to aphids by the method of paper partition chromatography. *Can. Entomologist*, 82 : 175-176. 1950.
2. BASU, A. C. Effect of different foods on the larvae and postlarval development of the moth *Prodenia litura* Fb. *J. Bombay Nat. Hist. Soc.* 44 : 275-288. 1943.
3. CHAUVIN, R. Influence du régime alimentaire sur le criquet pèlerin grégaire. *Compt. rend. soc. biol.* 131 : 31-33. 1939.
4. CREIGHTON, J. T. Factors influencing insect abundance. *J. Econ. Entomol.* 31 : 735-739. 1938.
5. DAHMS, R. G. Effect of different varieties and ages of sorghum on the biology of the chinch bug. *J. Agr. Research*, 76 : 271-288. 1948.
6. DAVIDSON, J. Biological studies of *Aphis rumicis* Linn. Reproduction on varieties of *Vicia faba*. *Ann. Applied Biol.* 9 : 135-145. 1922.
7. DAVIDSON, J. Biological studies of *Aphis rumicis* Linn. The penetration of plant tissues and the sources of food supply of aphids. *Ann. Applied Biol.* 10 : 35-54. 1923.
8. EVANS, A. C. Physiological relationships between insects and their host plants. I. The effect of the chemical composition of the plant on reproduction and production of winged forms in *Brevicoryne brassicae* L. (Aphididae). *Ann. Applied Biol.* 25 : 558-572. 1938.
9. HASEMAN, L. Influence of soil minerals on insects. *J. Econ. Entomol.* 39 : 8-11. 1946.
10. HILL, R. E. Influence of food plants on fecundity, larval development and abundance of the tuber flea beetle in Nebraska. *Nebraska Agr. Expt. Sta. Research Bull.* 143. 1946.
11. HODGE, C. Growth and nutrition of *Melanoplus differentialis* Thomas (Orthoptera: Acrididae). I. Growth on a satisfactory mixed diet and on diets of single food plants. *Physiol. Zool.* 6 : 306-328. 1933.
12. LEES, A. H. Insect attack and the internal condition of the plant. *Ann. Applied Biol.* 13 : 506-515. 1926.
13. LOTT, T. B. A mechanism for the automatic irrigation of sand cultures. *Science*, 88 : 17-18. 1938.
14. MA, T. S. and ZUAZAGA, G. Micro-Kjeldahl determination of nitrogen. *Ind. Eng. Chem. Anal. Ed.* 14 : 280-281. 1942.
15. MARTIN, H. The scientific principles of plant protection with special reference to chemical control. Chap. 2, 3rd ed. Edward Arnold and Co., London. 1940.
16. MUMFORD, E. P. and HEY, D. H. The water balance of plants as a factor in their resistance to insect pests. *Nature*, 125 : 411-412. 1930.
17. PEPPER, J. H. and HASTINGS, E. Biochemical studies on the sugar beet webworm (*Loxostege sticticalis* L.) with special reference to the fatty acids and their relation to diapause and sterility. *Montana Agr. Expt. Sta. Tech. Bull.* 413. 1943.
18. SEAMANS, H. L. and McMILLAN, E. The effect of food plants on the development of the pale western cutworm (*Agrotis orthogonia* Morr.). *J. Econ. Entomol.* 28 : 421-425. 1935.
19. SMITH, D. S., HANDFORD, R. H., and CHEFURKA, W. Some effects of various food plants on *Melanoplus mexicanus mexicanus* (Sauss.) (Orthoptera: Acrididae). To be published.
20. TAUBER, O. E., DRAKE, C. J., and DECKER, G. C. Effects of different food plants on egg production and adult survival of the grasshopper *Melanoplus bivittatus* (Say). *Iowa State Coll. J. Sci.* 19 : 343-359. 1945.
21. WEBSTER, J. E., SIEGLINGER, J., and DAVIES, F. Chemical composition of sorghum plants at various stages of growth, and relation of composition to chinch bug injury. *Oklahoma Agr. Expt. Sta. Tech. Bull.* T-30. 1938.



CANADIAN JOURNAL OF ZOOLOGY

Notes to Contributors

GENERAL: Manuscripts should be typewritten, double spaced, and the **original and one extra copy** submitted. Style, arrangement, spelling, and abbreviations should conform to the usage of this Journal. Names of all simple compounds, rather than their formulas, should be used in the text. Greek letters or unusual signs should be written plainly or explained by marginal notes. Superscripts and subscripts must be legible and carefully placed. Manuscripts should be carefully checked before being submitted, to reduce the need for changes after the type has been set. If authors require changes to be made after the type is set, they will be charged for changes that are considered to be excessive. **All pages, whether text, figures, or tables, should be numbered.**

ABSTRACT: An abstract of not more than about 200 words, indicating the scope of the work and the principal findings, is required.

ILLUSTRATIONS:

(i) **Line Drawings:** All lines should be of sufficient thickness to reproduce well. Drawings should be carefully made with India ink on white drawing paper, blue tracing linen, or co-ordinate paper ruled in blue only; any co-ordinate lines that are to appear in the reproduction should be ruled in black ink. Paper ruled in green, yellow, or red should not be used unless it is desired to have all the co-ordinate lines show. Lettering and numerals should be neatly done in India ink preferably with a stencil (do not use typewriting) and be of such size that they will be legible and not less than one millimeter in height when reproduced in a cut three inches wide. All experimental points should be carefully drawn with instruments. Illustrations need not be more than two or three times the size of the desired reproduction, but the ratio of height to width should conform with that of the type page. **The original drawings and one set of small but clear photographic copies are to be submitted.**

(ii) **Photographs:** Prints should be made on glossy paper, with strong contrasts; they should be trimmed to remove all extraneous material so that essential features only are shown. Photographs should be submitted in duplicate; if they are to be reproduced in groups, one set should be so arranged and mounted on cardboard with rubber cement; the duplicate set should be unmounted.

(iii) **General:** The author's name, title of paper, and figure number should be written in the lower left hand corner (outside the illustration proper) of the sheets on which the illustrations appear. Captions should not be written on the illustrations, but typed on a separate paper of the manuscript. All figures (including each figure of the plates) should be numbered consecutively from 1 up (arabic numerals). Each figure should be referred to in the text. If authors desire to alter a cut, they will be charged for the new cut.

TABLES: Each table should be typed on a separate sheet. Titles should be given for all tables, which should be numbered in Roman numerals. Column heads should be brief and textual matter in tables confined to a minimum. Each table should be referred to in the text.

REFERENCES: These should be listed alphabetically by authors' names, numbered in that order, and placed at the end of the paper. The form of literature citation should be that used in this Journal. Titles of papers and inclusive page numbers of the references cited should be given. All citations should be checked with the original articles. Each citation should be referred to in the text by means of the key number.

REPRINTS: A total of 50 reprints of each paper, without covers, are supplied free to the authors. Additional reprints will be supplied according to a prescribed schedule of charges. On request, covers can be supplied at cost.

Approximate charges for reprints may be calculated from the number of printed pages, obtained by multiplying by 0.6 the number of manuscript pages (double-spaced typewritten sheets, 8½ in. × 11 in.) and making allowance for space occupied by line drawings and halftones (not inserts). The cost per page is tabulated at the back of the reprint request form sent with the galley.

Manuscripts should be addressed: *Canadian Journal of Zoology,*
National Research Council,
Ottawa, Canada.

Contents

	Page
The Effects on the Grasshopper, <i>Melanoplus mexicanus mexicanus</i> (Sauss.) (Orthoptera: Acrididae) of Varying the Nitrogen Content in its Food Plant—D. S. Smith and F. E. Northcott - - - - -	297

25

